

Columns and supplies catalog



HPLC



GC



TLC




SPE and Flash



Syringe filters



Vials and caps



... we Meet your Needs

Since 1911 MACHEREY-NAGEL has represented high quality, innovation and reliability in chemical and biomolecular analysis. Friendly expert advice for our highly valued customers as well as outstanding product quality have been the cornerstones of our corporate success for more than 100 years. CEO C. Wagner, the great-granddaughter of the company's founder, has been managing the enterprise since 2000.

Our product ranges:



Filtration



Rapid Tests



Water Analysis



Chromatography



Bioanalysis

Dedicated from the very first:

Milestones of Chromatography at MACHEREY-NAGEL




1952:
MN launches the first products for paper chromatography. In the same year, Martin and Synge receive the Nobel Prize in Chemistry for the development of partition chromatography.



1961:
MACHEREY-NAGEL becomes one of the pioneers in TLC.



1970:
Expansion of the product range by column chromatography




1974:
NUCLEOSIL® · one of the first spherical HPLC silicas leads to our core competence in silica technology



1982:
first fused silica capillary columns for GC



1987:
CHROMABOND® columns for SPE



1994:
OPTIMA® capillary columns for optimum GC separations



2002:
NUCLEODUR® high purity spherical silica for HPLC

2011:
NUCLEOSHELL® core shell silica for highest efficiency in HPLC

Chromatography

Visit our Chromatography pages

www.mn-net.com

Chromatography



- Product information · always up-to-date
- ChromaNews
- Job opportunities
- Material safety data sheets
- Free application database with about **3000** chromatography applications



Solid Phase Extraction and Flash chromatography



CHROMABOND® phases and accessories for SPE	1-52
High-throughput SPE and Flash chromatography	53-64

Sample clarification



CHROMAFIL® syringe filters	66-76
----------------------------	-------

Vials and closures



Vials and closures	77-103
Autosampler compatibility	103-108

Liquid chromatography



Columns for HPLC	110-197
Packings for liquid chromatography	198-206

Thin layer chromatography



Plates, kits and adsorbents for TLC	208-234
-------------------------------------	---------

Gas chromatography



Capillary columns for GC	236-281
Reagents for GC	282-291
Accessories for GC	293-297

Appendices

Product and subject index, index of reference numbers, trademarks, USP lists	298-327
--	---------



New products for sample preparation

CHROMABOND® HR-Xpert


your way to cleaner samples by SPE

The professional concept of innovative RP and mixed-mode ion exchange phases for SPE

- ✦ CHROMABOND® HR-X · hydrophobic PS/DVB copolymer
- ✦ CHROMABOND® HR-XC and HR-XCW
strong and weak mixed-mode cation exchangers
- ✦ CHROMABOND® HR-XA and HR-XAW
strong and weak mixed-mode anion exchangers
- ✦ Now available in 2 particle sizes:
standard particle size 85 µm
particle size 45 µm for smaller sample volumes,
lower adsorbent weight and
smaller elution volumes


... Order your free
samples today!
info@mn-net.com

Page
10



New CHROMAFIL® filters

for optimum filtration results



- ✦ CHROMAFIL® Combi filters
Syringe filters with a coarse glass fiber prefilter
recommended for solution with a high particle load
- ✦ New dimensions and pore sizes for Nylon, cellulose acetate and PTFE filters

Page
68

Vials and closures

completely revised product range



- ✦ Large selection of
 - Crimp neck vials, crimp closures, crimping tools
 - Screw neck vials and screw closures
 - Special vials and closures
- ✦ High quality products at an equitable
price performance ration


Page
77

We look forward to sharing our expertise with you:
vials@mn-net.com



NUCLEOSHELL® core-shell silica

for rapid, efficient HPLC



- Solid core of silicon dioxide, homogeneous shell of porous silica
- Highest efficiency compared to totally porous materials
- Pore size 90 Å; particle size 2.7 µm (core 1.7 µm); specific surface 130 m²/g
- Lower back pressure enables use on conventional LC systems
- Pressure stability 600 bar

Page
144

NUCLEOSHELL® RP 18


nonpolar high density phase

Key features:

- Suitable for LC/MS and HPLC at pH extremes (pH 1–11)
- Superior base deactivation, ideal for method development

Recommended application:

Overall sophisticated analytical separations, e.g., analgesics, anti-inflammatory drugs, antidepressants; herbicides; phytopharmaceuticals; immunosuppressants
USP L1



Page
148

NUCLEOSHELL® PFP

hydrophobic pentafluorophenyl phase


Key features:

- Hydrophobic phase with alternative selectivity in comparison to classical C₁₈ modifications
- Separation principle based on 4 retention mechanisms (polar interactions (H bonds), dipole-dipole interactions, π-π interactions, hydrophobic interactions)

Recommended application:

Aromatic and unsaturated compounds, phenols, halogenated compounds, isomers, polar compounds like pharmaceuticals, antibiotics; high retention of basic compounds


USP L43



Page
150

NUCLEOSHELL® HILIC

zwitterionic phase



Key features:

- Ideal for reproducible and stable chromatography of highly polar analytes
- Very short column equilibration times

Recommended application:

Hydrophilic compounds such as organic polar acids and bases, polar natural compounds, nucleosides, oligonucleotides, amino acids, peptides, water soluble vitamins

Page
152



New products for HPLC

NUCLEODUR® PolarTec

RP phase with embedded polar group



Key features:

- Excellent base deactivation
- Suitable for LC/MS and stable in 100% aqueous mobile phases
- Pronounced steric selectivity

Recommended application:

Exceptional selectivity for phenols and nitrogen containing compounds, polar compounds like basic pharmaceuticals, organic acids, pesticides, amino acids, water soluble vitamins, etc.

USP L1 and L60

Page
124

NUCLEODUR® PFP

hydrophobic pentafluorophenyl phase



Key features:


- Hydrophobic phase with alternative selectivity in comparison to classical C₁₈ modifications
- Separation principle based on 4 retention mechanisms (polar interactions (H bonds), dipole-dipole interactions, π - π interactions, hydrophobic interactions)
- Suitable for LC/MS due to low bleeding characteristics

Recommended application:

Aromatic and unsaturated compounds, phenols, halogenated compounds, isomers, polar compounds like pharmaceuticals, antibiotics; high retention of basic compounds

USP L43

Page
136




ALUGRAM® Xtra

optimized TLC silica layers on aluminium

- ✦ Outstanding wettability for precise colorization results, even with 100% aqueous derivatization and visualization reagents
- ✦ Excellent separation efficiency and reproducibility from lot to lot
- ✦ Easy and reliable cutting due to an optimized binder system, no flaking of silica; now available as
 - ALUGRAM® Xtra SIL G with unmodified standard silica
 - ALUGRAM® Xtra Nano-SIL with unmodified HPTLC silica
 - ALUGRAM® Xtra SILGUR standard silica with concentrating zone
 - ALUGRAM® Xtra Nano-SILGUR HPTLC with concentrating zone




Pages
214-218



OPTIMA WAXplus® OPTIMA® FFAPplus

optimized polar phases for GC


- ✦ WAXplus
- ✦ Polar phase with improved cross-linking for reduced column bleeding and better temperature stability



- ✦ Good solvent stability, extended column lifetime, better performance in the MS in comparison to conventional Wax phases
- ✦ Recommended application: broad ranges of analyses, e.g., for solvents and alcohols, suitable for aqueous samples
- ✦ USP G16

Page
261

- ✦ FFAPplus
- ✦ Polar polyethylene glycol phase with nitro-terephthalic acid modification and innovative cross-linking



- ✦ Good solvent stability, extended column lifetime, better performance in the MS in comparison to conventional Wax phases, suitable for aqueous samples
- ✦ Recommended application: FAMES, allows determination of free carboxylic acids without derivatization
- ✦ USP G35 / close equivalent to G25

Page
263